

Absorbing Phase Transitions with Coupling to a Static Field and a Conservation Law

Julien Kockelkoren and Hugues Chaté
 CEA – Service de Physique de l’État Condensé,
 Centre d’Études de Saclay, 91191 Gif-sur-Yvette, France
 (Dated: February 2, 2008)

The coupling of branching-annihilating random walks to a static field with a local conservation law is shown to change the scaling properties of their phase transitions to absorbing states. In particular, we find that directed-percolation-class transitions give rise to a new universality class distinct from that characterizing the depinning of the so-called linear interface model.

PACS numbers: 64.60.Cn, 05.70.Ln, 82.20.-w, 89.75.Da

Among the many works aiming at an understanding of universality in out-of-equilibrium critical phenomena, those on transitions to absorbing states play a leading role because these phase transitions have no equilibrium counterparts and even occur in one-dimensional (1D) systems [1, 2]. In the simple and general case of the reaction and diffusion of identical particles A without site occupation restriction, recent numerical progress (awaiting analytical confirmation) has led to a global picture involving four basic universality classes [3]. Consider reactions of the type $[mA \rightarrow (m+k)A, nA \rightarrow (n-l)A]$ where m, n, k , and l are positive integers. Outside the prominent directed percolation (DP) class, whose simplest representatives are given by single-particle reactions ($m, n = 1$), the PCPD and TCPD classes (for “pair/triplet contact process with diffusion” — a vocable largely used for historical reasons) are respectively characterized by reactions involving two ($m, n = 2$) and three ($m, n = 3$) particles [4]. The fourth class has been considered to be defined by the conservation of the parity of the number of particles, hence its usual name PC (for “parity-conserving”) [1], but it is now clear that this is not its defining feature [5]. Nevertheless, the relevance of conservation laws is ascertained at equilibrium, and it remains important to explore it within absorbing phase transitions (APT), since conserved quantities abound in physical situations.

As a matter of fact, it was argued that DP-class problems where the order parameter is coupled to an auxiliary (diffusive) field with conservation generally show non-DP critical properties [6]. The particular case where the auxiliary field is static was conjectured recently to lead to yet another class of APT [7, 8]. (Note that the corresponding models then possess infinitely-many non-connected absorbing states and strong memory effects.) Numerical simulations confirmed only partially the above ideas. The situation, in our view, remains unsatisfactory for the static case: while in 1D no definite conclusions could be reached [9], in 2D and 3D the numerically-estimated critical exponents, when self-consistent, were found to roughly coincide with those of the depinning transition of the so-called linear interface model (LIM) [10]. A significant departure from DP-scaling was thus found, but the existence of a separate universality class remains in

question. Meanwhile, no rigorous result is available, although a heuristic mapping on the dynamics of LIM was advocated in [11].

In this Letter, we investigate the relevance of conservation laws to APT within the more general framework of four “basic” universality classes briefly recalled above, but restricting ourselves to the coupling to a static auxiliary field: now the creation of active, diffusive particles A is conditioned to the presence of passive, static particles B in such a way that the total number of particles is locally conserved. We thus study the reactions $[mA + kB \rightarrow (m+k)A, nA \rightarrow (n-l)A + lB]$, and find evidence for the existence of new universality classes. We revisit some of the models studied in [7, 8, 9], but we mostly deal with two-species reaction-diffusion models derived from those introduced in [3]. We show the existence of a specific “C-DP” (“DP with conservation”) class distinct from both DP and LIM in space dimensions 1 to 3, and point at a crucial difference with the LIM which was overlooked in [11]. We also show that a specific C-PCPD class exists, while no significant departure from TCPD and PC scaling could be observed in the presence of the coupling to a static field with conservation.

Extending the notation introduced in [3], we encode the rules by the order of the branching and annihilation reactions using letters s, p, t (standing for singleton, pair, triplet) for $m, n = 1, 2, 3$, followed by integers k and l . For instance, reactions $[2A + B \rightarrow 3A, 2A \rightarrow 2B]$ are coded as “conserved PCPD rule” C-pp12. Our two-species bosonic models are updated in two parallel substeps: A particles move to one of their nearest neighbors (strong diffusion), then on-site reactions take place, involving the n_A and n_B particles present locally.

We first present our results on the implementation of the C-ss11 rule studied in [8]. In this case, the branching reaction $A + B \rightarrow 2A$ is performed for each of the n_A particles present with probability $p(n_B) = 1 - 1/2^{n_B}$, while the annihilation reaction $A \rightarrow B$ is performed every A particle with probability q . Note that since $p(0) = 0$, branching is indeed conditioned to the presence of B particles. Unlike in [8], where the total density of particles is used as a control parameter, here we vary q , which

TABLE I: Critical exponents of the C-DP class in 1D, 2D, and 3D. The corresponding values for DP and LIM are also given for reference. Values for the LIM class, as well as estimates of κ for usual DP models were measured by ourselves with system sizes and timescales of the same order as those used for the C-DP class.

d		δ	z	β	2κ
	DP	0.1596	1.58	0.2765	0.84(1)
1	C-DP	0.140(5)	1.55(3)	0.29(2)	0.86(1)
	LIM	0.125(5)	1.43(1)	0.25(2)	0.35(1)
	DP	0.451	1.76	0.584	0.56(2)
2	C-DP	0.51(1)	1.55(3)	0.64(2)	0.50(2)
	LIM	0.50(1)	1.55(2)	0.63(2)	0 ⁺
	DP	0.73	1.90	0.81	0.30(5)
3	C-DP	0.88(2)	1.73(5)	0.88(2)	< 0.2
	LIM	0.77(2)	1.78(7)	0.85(2)	0

allows to start from perfectly homogeneous initial conditions (typically one A particle on each site). At large q , annihilation dominates and a static configuration of B particles is quickly reached, while at small q frequent branching ensures a stationary density of A particles.

Our numerical methodology is standard: monitoring various order parameters (e.g. ρ_A the density of A particles), we first determine the critical point q_c separating decay to an absorbing state from sustained activity, expecting then an algebraic law ($\rho_A \sim t^{-\delta}$) with $\delta = \beta/\nu_{\parallel}$. We record also, during these runs, the mean squared local gradient $\langle(\nabla h)^2\rangle$ of the interface $h(x, t)$, where $h(x, t)$ is the time integral of the local activity at site x until time t ($h \rightarrow h + 1$ whenever $n_A > 0$, with $h = 0$ initially). First introduced in [12] for DP-class models, this fictitious interface was shown to have “anomalous scaling” in the form of local gradients diverging algebraically at the critical point: $\langle(\nabla h)^2\rangle \sim t^{2\kappa}$. (Note that it is also the interface conjectured to behave like in the LIM by Alava and Muñoz for C-DP critical points [11].) After this first series of runs, we estimate the decay of the (stationary) order parameter with the distance to the previously-estimated threshold, yielding an estimate of exponent β . Finally, exponent z is estimated via the finite-size scaling of the lifetime of activity at threshold.

Table I summarizes our findings for the C-ss11 rule defined above in space dimensions 1 to 3. Since it is impossible, here, to present all the data which led to this table, we only show the most significant results, while a full report will appear elsewhere [13]. Two conclusions can be drawn. Firstly, we confirm that the coupling to the static particles B together with the conservation law does break the DP class. Our estimates in 2D and 3D are consistent with those published earlier [7, 8, 10], while in 1D (Fig. 1), our data exhibit more than 3 decades of scaling, leading to our estimate $\delta_{1D}^{C-DP} = 0.14(1) \neq \delta_{1D}^{DP} = 0.1596$




FIG. 1: 1D data from the C-ss11 rule implemented as in [8] and from the Leschhorn automaton for the LIM [15]. Near-critical decay from fully active homogeneous initial conditions (system size 2^{22-26} sites). (a): density of active particles ρ vs time (top 3 curves: Leschhorn model with $p = 0.80085, 0.8008, 0.80075$; bottom 5 curves: C-ss11 model with $1 - q = 0.82858, 0.8286, 0.82861, 0.82863, 0.82865$). Top inset: $\rho \times t^{0.14}$ vs time. Bottom inset: $\rho \times t^{0.125}$ vs time. (b): growth of $y^2 = \langle(\nabla h)^2\rangle$ during the same runs at criticality for the C-ss11 model ($1 - q_c = 0.82861$, top curve) and for the Leschhorn automaton ($p_c = 0.8008$, bottom curve).

(runs showing an effective decay exponent close to the DP value eventually become clearly subcritical). Secondly, our results also lead to ruling out LIM-class scaling: our estimates of exponents δ , β and z for this C-ss11 rule cannot fully rule out LIM values (Table I): no discrepancy could be registered in 2D, while in 1D and 3D we record significant but small differences (insets of Figs. 1 and 2). On the other hand, it is clear from our results on the “roughness exponent” κ that there are fundamental differences between a LIM interface and the fictitious interface constructed at the C-ss11 critical points: in 1D, the LIM interface shows anomalous scaling, but κ is more than twice smaller than for the C-ss11 rule (Fig. 1b). In higher dimensions, LIM interfaces are *not* locally rough ($\kappa = 0$), i.e. there is surface tension (Fig. 2). Meanwhile, we observe, to numerical accuracy, $2\kappa = 1 - \delta \neq 0$ for the C-ss11 critical points, as observed for ordinary APT [14].

In order to assess the robustness of the above results, we also studied a number of other rules whose “uncoupled” version have APT in the DP-class, implemented in the spirit of [3]. Consider for instance rule C-sp12, encoding reactions ($A + B \rightarrow 2A$, $2A \rightarrow 2B$). For each of the $\lfloor n_A/2 \rfloor$ pairs of active particles present locally, either the branching or the annihilating reaction takes place with probability p or $q = 1 - p$. Here p is independent of n_B , but the branching reaction is further subjected to the presence of enough B particles among those present initially. Finally, if $n_A = 1$, only the branching reaction can occur (with probability p and if $n_B > 0$).

Rule C-ss11, implemented as above, was investigated in




FIG. 2: Same as Fig. 1, but in 2D and 3D Top left: near-critical decay of ρ in 2D for C-ss11 rule (top curve, $1 - q = 0.65298$, $2^{14 \times 2}$ sites) and Leschhorn automaton (bottom curve, $p = 0.64173$, $2^{15 \times 2}$ sites); Top right: same in 3D for C-ss11 rule (bottom curves, $1 - q = 0.57908$ and 0.57909 , $2^{11 \times 3}$ sites) and Leschhorn automaton (top curves, $p = 0.52722$ and 0.527225 , $2^{10 \times 3}$ sites); Bottom panels: $y^2 = \langle (\nabla h)^2 \rangle$ for the same runs (left: C-ss11 rule, right: Leschhorn automaton).




FIG. 3: Same as Fig. 1a, but for C-DP rules implemented following [3] (see text). Only near-criticality runs are shown. From top to bottom: C-ss11 rule ($p = 0.720775$, 2^{26} sites); C-ss21 rule ($p = 0.75905$, 0.75906 , 2^{24} sites); C-sp12 rule ($p = 0.7094$, 2^{22} sites); C-ps11 rule ($p = 0.83811$, 2^{24} sites). Inset: $\rho \times t^{0.14}$ vs time for the same data.

1, 2, and 3 dimensions, yielding exponent values equal, to numerical accuracy, to those reported in Table I. Rules C-ss21, C-sp12, and C-ps11 were also studied thoroughly in 1D with, again, exponent values fully compatible with those found for rules C-ss11 (Fig. 3). All these results give credence to the existence of specific C-DP universality




FIG. 4: Same as Fig. 3, but for C-PCPD rules, for systems of size 2^{24} or 2^{26} sites. Top 5 curves: C-pp12 rule (from bottom up: $p = 0.81813$, 0.81814 , 0.81815 , 0.81816 , 0.81817). Bottom 5 curves: C-pp22 rule (from bottom up: $p = 0.82305$, 0.82308 , 0.82309 , 0.8231 , 0.82312). Top inset: $\rho \times t^{0.17}$ vs time. Bottom inset: $\rho \times t^{0.2}$ vs time.

class, distinct from both the DP and the LIM classes.

We now report on our results on conserved PCPD, TCPD, and PC systems, which we only investigated in 1D. For rules C-pp12 and C-pp22, our estimates of critical exponents reveal an identical departure from the PCPD class which we interpret as testifying to the existence of a C-PCPD class. In particular, the critical decay experiments allow to clearly rule out the PCPD value of the δ exponent (Fig. 4). We find, for both rules studied: $\delta = 0.17(1)$, $\beta = 0.32(2)$ and $z = 1.55(5)$, whereas our estimates for the PCPD class are $\delta = 0.20(1)$, $\beta = 0.37(1)$ and $z = 1.70(5)$. The careful study of rules C-tt12 and C-tt22 did not reveal any significant departure from TCPD scaling for these C-TCPD rules. Similarly, we found only small influence of the coupling to a static field for generalized voter (PC) rules C-mp22 and C-mp42 [13]. We have, at present, no reason to believe that the C-PC and C-TCPD classes “do not exist”.

Before our main conclusions, we come back to the suggested equivalence between the LIM and C-DP classes which is contradicted by our numerical results. In the Leschhorn automaton [15], used here for our estimates of LIM class exponents, an integer-height interface evolves in parallel according to:

$$h_i \rightarrow h_i + 1 \quad \text{iff} \quad \nabla^2 h_i + f_i > 0 \quad (1)$$

where i is a lattice index and $\nabla^2 h_i$ is the discrete Laplacian calculated with the nearest-neighbors of i . A key ingredient is the quenched bimodal random “force” f_i ($f_i = \pm 1$ with probability p and $1 - p$) which is redrawn every time the interface advances.

The evolution of the fictitious interfaces from which exponent κ is estimated in C-DP models can be cast into an equation very similar to (1) [11]. This is perhaps best seen when considering the Manna or “fixed energy”

sandpile, a much-studied model of the C-DP type which, unfortunately, suffers from severe corrections to scaling [9, 13, 16]. There, if the number z_i of “sand grains” present at lattice site i is equal to or exceeds a threshold z_c usually taken to be 2, then z_c grains independently hop to a randomly-chosen nearest-neighbor. This defines active and passive sites, corresponding to the above A and B particles, and the local conservation law is obviously that of the sand particles. The fictitious interface thus simply encodes the number of times a site has discharged its grains since $t = 0$: $h_i \rightarrow h_i + 1$ when $z_i \geq z_c$. This local discharge condition can be expressed in terms of z_i^{in} and z_i^{out} , the cumulated number of particles received from and distributed to the neighbors. Consider, for simplicity, the 1D case with $z_c = 2$. Clearly, $z_i^{\text{out}} = 2h_i$, whereas z_i^{in} can be written as the sum of a mean flux ($h_{i+1} + h_{i-1}$) and a fluctuating part $\tau_i(h_i)$ [11, 17]. The discharge condition is then expressed

$$h_{i+1} + h_{i-1} - 2h_i - 1 + z_i(0) + \tau_i(h_i) > 0$$

where $z_i(0)$ is the initial number of particles on site i . In the general case, the fictitious interface thus obeys

$$h_i \rightarrow h_i + 1 \quad \text{iff} \quad \nabla^2 h_i - 1 + z_i(0) + \tau_i(h_i) > 0 \quad (2)$$

which is, up to the presumably irrelevant “columnar noise” $z_i(0) - 1$, formally equivalent to Eq.(1). There exists, however, a fundamental difference between the two equations: whereas f_i is a delta-correlated noise, τ_i is a sum of random variables and its amplitude diverges as $t \rightarrow \infty$. As a matter of fact, since z_i is bounded, both $\langle(\nabla^2 h)^2\rangle$ and $\langle\tau(h)^2\rangle$ must diverge in the same way. We have checked numerically that it is the case, and found moreover that $\langle(\nabla^2 h)^2\rangle \sim t^{2\kappa}$. On the other hand, for the Leschhorn automaton interfaces, the amplitudes of both the noise and the Laplacian remain bounded, indicating that it constitutes a reasonable physical interface model, even though $\langle(\nabla h)^2\rangle$ diverges in 1D.

These results indicate that the connection between C-DP and LIM via the fictitious interfaces is only formal and bears no reality. Thus, the numerical values of exponents δ , z and β are probably *coincidentally* very close to each other in both classes, and C-DP critical scaling defines a universality class distinct from that of the LIM.

To sum up, we have found evidence that the coupling of usual reaction-diffusion systems to a static field with a local conservation law changes the scaling laws of their transitions to absorbing states, giving rise to bona fide universality classes distinct from previously known ones. Naturally, these results call for rigorous analytical confirmation which will probably only happen when the classification of simple APT (DP, PCPD, etc.) is itself given some solid theoretical basis.

Reaching a global understanding of absorbing phase transitions with coupling to an auxiliary field requires, from our viewpoint, to study the respective importance

of the static or diffusive character of this field and of the presence of a conservation law. In particular, the sole coupling to a static field (without conservation) is usually believed to be irrelevant from existing work on simple fermionic models such as the pair contact process [18]. This should now be tested against the general framework put forward here. Similarly, the case of a diffusive field with conservation should be extended beyond the DP-class. Both directions are currently being investigated.

We thank M.A. Muñoz and F. van Wijland for many fruitful discussions.

-
- [1] H. Hinrichsen, Adv. Phys. **49**, 815 (2000).
 - [2] G. Ódor, eprint cond-mat/0205644, and references therein.
 - [3] J. Kockelkoren and H. Chaté, Phys. Rev. Lett. **90**, 125701 (2003).
 - [4] For a recent account of an ongoing debate, see G. Ódor, eprint cond-mat/0304023, to appear in Braz. J. Phys.
 - [5] Within our framework, the *only* rules in the “parity-conserving” class are $[A \rightarrow (1+k)A, 2A \rightarrow \emptyset]$ with k even, i.e. those which can be mapped onto models with two symmetric absorbing states, suggesting that the critical behavior is that of the (generalized) voter model (see I. Dornic, H. Chaté, J. Chave, and H. Hinrichsen, Phys. Rev. Lett. **87**, 045701 (2001)).
 - [6] R. Kree, B. Schaub, and B. Schmittmann, Phys. Rev. A **39**, 2214 (1989); F. van Wijland, K. Oerding, and H.J. Hilhorst, Physica A **251**, 179 (1998); K. Oerding, F. van Wijland, J.-P. Leroy, and H.J. Hilhorst, J. Stat. Phys. **99**, 1365 (2000).
 - [7] M. Rossi, R. Pastor-Satorras, and A. Vespignani, Phys. Rev. Lett. **85**, 1803 (2000); A. Vespignani, R. Dickman, M.A. Muñoz, and S. Zapperi, Phys. Rev. E **62**, 4564 (2000).
 - [8] R. Pastor-Satorras and A. Vespignani, Phys. Rev. E **62**, R5875 (2000).
 - [9] R. Dickman et al., Phys. Rev. E **64**, 056104 (2001).
 - [10] S. Lübeck, Phys. Rev. E **64**, 016123 (2001); **66**, 046114 (2002).
 - [11] M. Alava and M.A. Muñoz, Phys. Rev. E **65**, 026145 (2002).
 - [12] R. Dickman and M.A. Muñoz, Phys. Rev. E **62**, 7632 (2000).
 - [13] J. Kockelkoren and H. Chaté, to be published.
 - [14] J. Kockelkoren et al., in preparation.
 - [15] H. Leschhorn, Physica A **195**, 324 (1993); M. Jost and K.D. Usadel, Physica A **255**, 15 (1998).
 - [16] R. Dickman and J.M.M. Campelo, eprint cond-mat/0301054; R. Dickman, eprint cond-mat/0209589.
 - [17] M. Paczuski, S. Maslov, and P. Bak, Phys. Rev. E **53**, 414 (1996); M.J. Alava and K.B. Lauritsen, Europhys. Lett. **53**, 563 (2001).
 - [18] I. Jensen and R. Dickman, Phys. Rev. E **48**, 1710 (1993); R. Dickman, Phys. Rev. E **53**, 2223 (1996); M.A. Muñoz, G. Grinstein, R. Dickman, and R. Livi, Phys. Rev. Lett. **76**, 451 (1996); J. Stat. Phys. **91**, 541 (1998).